



## Don't forget the annual flu shot!

- -Recommended for **everyone ages 6 months or older**, if eligible *(see CDC Flu website)*.
- -The flu shot is safe to receive concurrently with a COVID-19 vaccine or booster dose.

## **Primary vaccination series**

Vaccine	Age	Dosing
Pfizer- BioNTech (mRNA)	6 months - 4 years  5 years and older	Three doses. Dose 2: at least 21 days after primary dose.  Dose 3: at least 8 weeks after second dose.   Two doses. Dose 2: at least 21 days after primary dose.
Moderna (mRNA)	6 months and older	Two doses. Dose 2: at least 28 days after primary dose.
Novavax (protein subunit)	12 years and older	Two doses. Dose 2: at least 21 days after primary dose.
Johnson & Johnson (viral vector)	18 years and older	One dose. Only recommended for adults unable to receive any other COVID-19 vaccine type.

## **Booster shot**

Recent COVID-19 infection? Wait three months after positive test date or start of symptoms to receive booster dose if due.

n	Bivalent* mRNA Booster (Pfizer or Moderna)	5 years and older **	Administer at least two months after finishing primary vaccine series or last booster dose.
	Novavax Booster	18 years and older	Administer at least six months after finishing primary vaccine series ONLY IF no previous booster dose(s) received.

<sup>\*</sup> Booster shots have been updated to include Omicron BA.4 and BA.5 subvariants.

<sup>\*\* 5</sup> year olds should receive a Pfizer Booster. People 6 years of age and older may receive either bivalent mRNA booster.



## More Information:

CDC: Influenza (Flu). https://www.cdc.gov/flu/index.htm

OHA: Vaccines, Boosters, and Third Doses. https://govstatus.egov.com/or-oha-booster-covid-19-vaccine

CDC: Stay Up to Date with Vaccines. https://www.cdc.gov/coronavirus/2019-ncov/vaccines/stay-up-to-date.html

CDC: COVID-19 Vaccine. https://www.cdc.gov/vaccines/covid-19/downloads/COVID-19-immunization-schedule-ages-6months-older.pdf

